JELLIUSSN 2519-5789 ONLINE ISSN:2320-3145

SCHOLARS International Refereed Multidisciplinary Journal of Contemporary Research

VOLUME III

(SPECIAL ISSUE)

FEBRUARY- 2015

IMPACT FACTORS

Universal Impact Factor = 1.2242 International Impact Factor Services = 0.654 Scientific Journal Impact Factor = 3.552 Global Impact Factor = 0.311

INDEXING

Directory Of Open Access Journals, Sweden Ulrich's Web-global Series Directory, USA Open J-Gate, India Advanced Science Index (ASI), Germany Cite Factor- Academic Scientific Journal- Canada/USA Academic.edu - Share Research- USA Academic Keys- Unlocking Academic Careers Yampu - Switzerland .docstock- We make your business better- Santa Monica, CA DRJI- Directory of Research Journals Indexing, India BASE- Bielefeld Academic Search Engine Calameo- Publish, Share, Browse- USA Indian Citation Index- India Slide Share- News Letters- San Francisco Scientific Indexing Services worldcat.org Pub-Res- (International Research Library)



MAAZ PUBLICATIONS

www.scholarsworld.net

:0

39

96

WATERSHED MAP FOR RIVER WATER POLLUTION CONTROL AND PREVENTION: WITH SPECIAL REFERENCE TO SELECTIVE PART OF GODAVARI RIVER IN NASHIK	97-99
AIR BORNE MYCOFLORA OF EXTRA AND INTRAMURAL ENVIRONMENT OF COLLEGE LIBRARY PATEL S.	100-104
A GEOGRAPHICAL STUDY OF POPULATION AND RESOURCES IN THE NASHIK DISTRICT PRALHAD Y. VYALIA	105-109
SOIL ANALYSIS OF THREE DIFFERENT LOCALITIES AROUND MALEGAON CITY KAPADNIS SHITAL SANTOSH	110-111
"IMPACT OF CLIMATIC CHANGE ON CROPPING PATTERN A CASE STUDY OF MALEGAON TAHSIL IN NASHIK DISTRICT" (MAHARASHTRA) SAWANT DINKAR DAULAT, SURYAWANSHI UTTAM PANDIT	经证据 医甲基甲基甲基
ECOTOURISM POTENTIAL OF NASHIK DISTRICT: A CASE STUDY OF BRAHMAGIRI AND SALHER FORT RAKESH V. PATIL, R. S. DEORE	116-120
32. ENVIRONMENTAL POLLUTION, PROBLEMS AND MITIGATION P.S. KUDNAR, C.M.NIKAM	121-125
LAND USE AND LAND DEGRADATION OF MALEGAON TAHASIL IN NASIK DISTRICT (MS): A GEOGRAPHICAL PERSPECTIVE U.P.SURYAWANSHI, D.D.SAVANT	126-129
TRIPLE ACTIVITY NATURAL ECOFRIENDLY ORGANOPHOSPHORUS (MONOCROTOPHOS) DEGRADING BACTERIAL ISOLATE FOR CLEANUP AND ENRICHMENT OF SOIL SEEMA GHARATE, RATAN BORSE, CHANDRAKANT SASLE	130-134
THE ADVERSE EFFECTS OF NON-SEASONAL PRECIPITATION ON AGRICULTURAL PRODUCTION IN DHULE DISTRICT, MAHARASHTRA PRIYANKA NIKUMBH, D. S. KUMBHAR	135-135
R. S. NIRWAN, A. B. SAWANT	136-138
POTENTIAL OF WIND ENERGY IN SURGANA TEHSIL B.A.AVHAD, J.V.MISAR	139-141
ONE NEW SPECIES OF THE GENUS TREMIORCHIS, MEHRA AND NEGI, 1926, AS TREMIORCHIS SPINENSIS IS COLLECTED FROM RANA TIGRINA @ SONGIR DIST DHULE M. S. INDIA D. N. PATIL, L. B. PAWAR	42-144

AIR BORNE MYCOFLORA OF EXTRA AND INTRAMURAL ENVIRONMENT OF COLLEGE LIBRARY

PATEL S.I.

Arts, Science and Commerce College, Ozar (MIG), Nashik-Maharashtra, India

ABSTRACT

The present investigation was undertaken to find out the concentration of the various air borne fungal spores in the outdoor and indoor environment of the College library. The mycoflora of indoor and outdoor environment of library in Ozar (MIG) College, Dist. Nashik (M.S) was investigated for the period of one year i.e. from October 2013 to September 2014 by Himedia Petri plate air sampler No. LA-030 with Himedia readymade Rose Bengal Chloramphinicol Agar culture plates techniques. There were total 21 air borne fungal colonies were isolated and identified in the ambient air of college library. Aspergillus (19.82% and 28.2%), Cladosporium (8.99% and 7.20%) Penicillium 8.23% and 5.30%), Fusarium (5.34 % and 4.50%) and Alternaria (6.10 % and 9.50%) were found to be all time occurrence and shown predominance in the outside and inside the library environment respectively. Lignocellulosic degrading fungus like Chaetomium (5.18% and 4.19 %) was also recorded in high humid and low temperature conditions, Alternaria, Fusarium Helminthosporium, Trichoderma, Rhizopus, Curvularia, Tetraploa, Periconia, Stemphyllum and some unidentified colonies were recorded in high concentration during the investigation with little variation in their concentration. The Total Colony Forming Units per M3 of air was more inside the library (1782/M3 of air) than outside (1680/M3 of air). The air borne fungal spores recorded in the library environment would be the main cause of some health hazards like sneezing, watery eyes, coughing and uneasy feelings in some sensitive visitors of the library. The air borne fungal spore concentration also shows close relation with the meteorological parameters like temperature, humidity and rainfall.

Keywords: Library, Allergy, Airborne Fungi, Weather Conditions

INTRODUCTION:

Due to suspended air borne particles in the library environment, both the readers and books get infected at more or less percentage. The enclosed environment of the library offer unique substrate such as binding glue, paper and other organic material which support the active fungal growth. Because of light weight and microscopic nature, the spores of fungi remain suspended in the air for long period. The concentration of these bioparticles shows great variation from time, season, altitude and weather conditions (Lacey 1991). Since the study of allergic aerospora of library environment is of great importance and continuous exposure to the library causes the respiratory disorders, A very meagre studies were reported on similar matter (Burge 1978, Vittal 1985, Tripathi 1987, Singh 1990, Agashe 1996, Ghani 1993, Verma 1996, Tilak 1997 Saoji 1997, Atluri 2002, Md. Shabbir 2003, Upadhyay 2005, Giri 2012,) hence the present work is undertaken to

study the allergic air borne fungal spores in the ambient air of library. Respiratory allergy is most prevalent among all populations with increasing trend all over the world. The bioparticulates implicated to cause allergic symptoms are pollen grains, fungal spores, insect debris, house dust mites, animal dander, chemicals and food etc. Today, more than 300 million of the world population is known to suffer from one or other allergic ailments affecting the socioeconomic quality of the life. Most people spend maximum of their time in the indoor environments of house, offices, factories and other indoor work places. This continuous exposure to the close indoor environment to many air borne pollutants comes through inhalation of air. The respiratory allergy is the most severe disorder including asthma, rhinitis, dermatitis, hay fever, skin rashes, sneezing, coughing, running nose, watering eyes etc. The major causative agents for the respiratory allergy are pollen grains, fungal spores, dust, mites, insect debris etc (Pande 2012).

MATERIAL AND METHOD:

The town Ozar (MIG), Maharashtra, is well known for its grape and onion production and similarly for air craft production at Hindustan Aeronautics Limited. The arts, Science and Commerce College, Ozar (MIG) was established along with central library in the year 1985. The total carpet area of the library is 1901 sq. fts consisting total 24,130 valuable books including text, reference books, periodicals and journals worth of Rs.27,38018/- up to year March 2014. The books were arranged subject wise and kept on open and closed steel racks keeping 3" distance between the racks.

The air sampling in the College library was done by using Himedia air sampler No. 030 with readymade Himedia Rose Bengal Chloramphinicol nutrient media plates. The air sampler was loaded with nutrient media plates and placed on the reading table for 10 minutes every forth night between 11.00 am to 11.30 am, i. e during peak hours of book transactions. The sampling of the air was done outside and inside the library environment from the months of October 2013 to March 2014in the first season and April to September 2014 in the second season. After the exposure, the petriplates are brought to the botany laboratory for the incubation at the temperature 25-27 °C for 4-5 days. The Colony Forming Units (CFU) were counted and Total Fungal Counts (TFC) were expressed as Colony Forming Units per cubic meter of air (CFUs/M3).

The fungal colonies detected per unit volume of air are calculated as under;

CFUs/M3= Number of Colonies on nutrient medium plates (N C) X 25 Sampling time in minus (T)

After the incubation period of 5-6 days the colonies were appeared on the nutrient medium petriplates. These colonies were counted and identified up to genus level with the help of available literature (Barnett1971, Tilak 1985).


RESULTS AND DISCUSSION:


Studies reveals the total 21 types of fungal colonies recorded in the ambient air of the library. High concentrations of fungal colonies were recorded in the library environment (326) than outside the library. Highest concentration of colonies was recorded to be of Aspergillus (inside 28.57 % and outside 17.485 %). Cladosporium (13.19% and 12.14 %), Penicillium (8.9 % and 4.29%), Helminthosporium (6.13 % and 5.0%) and Alternaria (4.29% and 7.86%) to the total colonies recorded inside and the outside the library environment respectively. (Table 01 and 02) In addition to the cellulose, the paper also contains the lignin, hemicellulose, pectin, waxes, tannins and minerals useful for the fungal growth. Beside the impurities formed during the paper production are helpful for the nutrition of fungi (Tilak S T 1985). The cellulytic activity of some paper deteriorating fungi produces endogluconase enzymes which damages the paper quality of the library book (Verma K S 1987) indicate the role of fungi in book deterioration.

Present studies indicates the high concentration of the fungal colonies outside the library in the months of October to March 2013 while there was gradual increase the concentration inside the library from the Months of April to September 2014 might be due to high humid and low temperature conditions. (Fig.01 & 02). Some fungal types like Drechslera, Epicoccum, and Geotrichum were only reported in the rainy months of July to September 2014. High concentration of fungal colonies (713) inside the library environment indicates humid and damp environmental conditions which would be favourable for the growth of moulds on the books. Chauhan et al (2006) reported the similar observation by studying 7 different libraries at Agra. The adverse effects of fungal components not only deteriorate the paper material but also affect the library staff. Agashe (1991) also reported 23 types of fungal colonies in the indoor environment of library at Bengaluru. Giri S K (2012) reported 36 fungal spore types from the ambient air of library at Nagpur adding highest concentration of Aspergillus, Cladosporium, Alternaria and Rhizopus spores. Predominence of Aspergillus and Cladosporium in the present studies is in relation with the reports of Vittal (1985) and Singh (1990). The cellulose degrading fungi like Chaetomium was also recorded (5.18% and 4.19 %) indicating the threat to the stored books in the library environment. Dhawan (2005) critically reviewed the role and aerobiodeteriogens on paper and museum material indicating high concentration of versatile fungi like Aspergillus, Penicillium, Cladosporium and Fusarium. As indicated by Nandimuthan (1995) and Tripathi (1987) the present studies also reveals high concentration of fungal colonies during the high humid, low temperature conditions in the month of December 2013 and January 2014 than in high temperature and low humid conditions in the month of February and March 2014. Since the study of allergic airospora of library environment is of great importance and continuous exposure to the library causes the respiratory disorders, A very meagre studies were reported on similar matter (Tilak S T (1982), Verma K S (1987), Tripathi S N (1987), Nandimuthan (1995), Shaney (2001), Sharma (2004), Tiwari K L (2004). And Mujumdar (2005). The studies inside and outside the library environment is of great importance since the library environment is loaded with the biopollutants that can be harmful to the health of the library users, beside reducing the life of books (Agashe S N 1996)

ACKNOWLEDGEMENT:

The author is extremely thankful to The Director, BCUD, Savitribai Phule Pune University, Pune for the financial assistance and The Principal Dr. A.P.Patil, MVP Samaj's Arts, Science and Commerce College, Ozar (MIG) for permitting air sampling in the College library and providing laboratory facilities.





578%

id los

mental

catal

dier el

diverse

staff

rary a

vaspur

pores

chous

bonded

tawan

cating

high nonth in the ment (S. A. 187), (04), reat if to

ine ind ing

TABLE: 01

					9	TAID	LE: ()1							
Fungal Types	E CONTRACTOR OF THE PARTY OF TH		Nur	nber o	of col	onies	recor	ded or	the					e library)	
Alternaria	3	Nov	Dec	Jan	Feb 1	Mar .	Apr A	Tax lar	un	etrip	lates	(Out	side th	e library)	
Aspergillus	12	4	1	21	1	3	-	No. 21000 8100	1	Tul.	Mug	Se	o Tot	al % Cor	ne CFU/I
Cercospora	3	2	211	12	7	8	6	8	A STATE OF THE PARTY OF	5	8	1 3	40	6.10	100
Chaetomium	3	5	2	-	-	2	-	- 1	Carlotte Balletin	37 24	22	-17	130	19.82	325
Cladosporium	12	3	4		STATE OF THE PERSON		3		the sales and	Harris I	5	-	17	2.59	42.5
Corynospora	1000		10	energy of the	Windson .		2 6	The second second second	L'analysis.	OR SEE S	3	3	34	5.18	85
Curvularia	-	SPECIAL CO	1,000	4			2		53.5		2	2 	59	8.99	147.5
Drechslera			SAMPLE OF THE PARTY OF THE PART	4	- 1 - 1	East Man	. 2	5	_	essed NS	5		17	.2,59	42.5
Epicoccum	2	1	- Commission	2 1					-		Maria Carl	of the second	20	3.05	50
Fusarium	3	2	4	man of the second			CIA LUNCO	-	3		7	-	17	2 5 S	
Geotrichum	2	1720	3 -	4	THE REAL PROPERTY.	CORPORATION POR	1	2	3 4		3	2	35	2.59	42.5
Helminthosporium	2	SCHOOL SECTION	6 4		6	18 91	-	-	-		-	-	15	5.34	87.5
Nigrospora	-	-	1 1	2	21/0941	0 100-3	3	1			3	3	130%	2.29	37.5
Penicillium Phoma	3	5	7 8		2	2		3	-		2	2	11	1.68	75
Renconia		16 -	-	-	10140	F186425	3	143.	7		7. 1	3	54	8.23	22.)
Pringsheimia	741 10	1 2			10000			- NUMBERS	-			-	1	0.15	135
O. C. Strategic Control of the Contr		- -	1	1	-	2 100 2 开线的	SHAPIG.	6	3		2	5	21	3.20	2.5 52.5
tan-1 11	200000	2	3	1	NAME OF THE OWNER, OWNE	100 to	-	- 10 A 3 A 3 A 3 A 3 A 3 A 3 A 3 A 3 A 3 A	1	-		2	5	0.76	12.5
The state of the s	Acres (7	2	1	PARK	1	3	3	1860		3	29	4.42	72.5
Inclassic 1	STIES HILLIANDS	2007.200	2	4	5	2	Table 1	2	6	-	To U.S. September	4	34	5.18	85
A STATE OF THE PARTY OF THE PAR	100	7	4	2	7	4	3	3	7	1	Seaton Supplies	2	29 1	4:42	72.5
Otal 57	/ 54	68	71	46	45	23	34	50	-	3	Service Commen)	58	8.84	145
				,	ГАВ	2012/28-85E-6		によりい。	63	82	6	3 (16	56	100	1640

72								E: 02		This will a			A SAN LAND	DO POL	04. 1.1.1640
Fungal Types	Oct	A Tool	N	umbe	er of	colon	ies re	corde	ed on t	he petr	:1				
Alternaria	2	Nov 3	Dec 2	Action to Beauty	Feb	Arten	Apr	May	Jun	Jul	Aug	N. T. Consultation of the	de the	library)	(1
Aspergillus	333	3	2	10		2	3	-	23	12	9	Se _E	-	al % Co	ne CEU/M
Cercospora	-	-	2	2	12	5	2	17	33	421	23	44	68	and the second	170
Chaetomium	(2000) (2000)			2			3	- Separate	5	4	2	7	27	3.79	THE PERSON OF TH
Cladosporium	2	4	5	3	2	1	9	7	7	5 E	17	3	30	4.21	67.5
Corynospora Curvularia	5.50	(5.15)	1071		with the second	ALE	13	2	2	5	7	4	51	7.15	127.5
Drechslera -	20020 59	- /	-	-	2	1	3	- Village	4	Sale in the	7		16	2.24	40
<i>Epicoccum</i>							婚者		3	5	2	7	24	3.37	60
Eusarium (1994)	2			2	-	-	2	-	1	4	-	2	12	1,68	1 30 B
Geotrichum	- 1	-	-	第一届	1788		4	2	4	8.4	7	493	9 32	1.26	22.5
Helminihosporium		100	2	1000	2		2	-	3	-	4	6	15	2.10	80
AND THE PARTY OF T	- J		2	1	1	- 特殊		3	3	5 (6)	3	8	28	3.93	37.5
Phoma	8.0	23	[編]				2		3	7	2	- 1	9	1.26	22.5
S Periconia		1781 1834		SEC LAN		- -	- 1000	1	-	7	3	6	38	5.33	95
Pringsheimia -	1		選 1891			1000	CONT. 1-6/20PM	2	123	4 1	2000	2	11	1.54	27.5
Rhizopus 2		1		- Maria	- NO 1866	2	1 .	-	-	1	- -	200		2.10	*37.5
Stemphyllum 2	1	3	1	2	2		1 2	1000		3	7	7		0.56	10
Trichoderma 2	1 1						-	The second		4	1 7	122000 1830	The second second	4.63 2.38	82.5
Unclassified 3 Total 21	4	2	1	1	4	3	3	14.7400	- The second	7	このでは からのと からかん	Total Control of the last	100000000000000000000000000000000000000	2.81	42.5
1 Lotal 21	22	28	26	25	man Property	THE RESERVE OF THE PARTY OF THE	Marie Committee of the			OF CHARLES STORY	2 6	5 5	-	7.43	132.5
						and and a second	THE PARTY	THE PERSON	1.11	13 11	7 12	1 7		00%	1782.5

REFERENCES:

- Agashe, S.N and Anuradha. (1996). Aeromycological studies of a library in Bangalore city, Ind. Journ. Aerobiol, Vol. 11: 24-28.
- Atluri, J.B and Padmini. (2002). Aeromycoflora of Andra university main library, Indian Journal of Aerobiology. Vol 18 (II), pp 47-58.
- 3. Barnett, H.L. and Hunter, B.B. (1972). Illustrated genera of imperfect fungi, Burgess. Co. Minneapolis, Minnesota, U.S.A.
- Burge H A .(1991). Bioaerosols: prevention and health effects in the indoor environment. Journ. Allergy clinical immunology, vol (26) 05, 687-701
- 5. Chauhan, S.V.S, A Kulkarni and Goyal .(2006). Air borne fungal spores of Agra city with special reference to allergic significance, *Indian Journal of Aerobiology*. Vol 17, pp 17-22.
- 6. Dhavan Sashi, and Neet Nigam .(2005).Fungal diversity of paper material and its role in bio deterioration in fungal diversity and biotechnology, M.K Rai and S.K. Deshmukh Scientific publisher Dodhpur, India PP 93-113.
- 7. Ghani, T and Shanta Kale .(1993). Aeromycoflora inside and outside the College library. *Indian Journal of Environ. Health*. Vol 3 (II), pp 1-5.
- 8. Lacey. J. (1991). Aerobiology and health. Role of aerobiology Fungal spores in respiratory diseases in, Oxford and IPH publishing Co. Ltd, New Delhi
- 9. Mohammad Shabbir, S. Nicky, Johnson and Rajasab. (2003). Aeromycoflora of Gujarat University library, *Indian Journal of Aerobiology*. Vol 16, pp 28-32.
- 10. Mujumdar M R and Surendra Hazare. (2005). Assessment of fungal contaminents in library of Kolkata, *Ind. Journ. Aerobiol.* 13: 398-403.
- 11. Nandimuthan and Vittal B P R. (1995). A volumetric survey of culturable moulds in Library, *Ind. Journ. Aerobiol.* 8:12-16.
- 12. Pande B.N. (2012). Bioaerosols as allergens, Bionano frontier, Vol. 5 (2), 164-169.
- Saoji A A and Giri S K .(1997). Concentration of aero allergenic fungal spores in intramural environment of Nagpur City-Hospital ward and library. 5th International conference on Aerobiology pp-211-218.
- 14. Shaney, Manju and Purwar. (2001). Aeromycoflora inside a library of Allahabad University, Ind. Journ. Aerobiol.14: 20-23.
- 15. Sharma K B and Basumaitrai. (2004). Air borne fungal spores in library and their association with paper and leather material, *Ind. Journ. Aerobiol.* 14: 35-43.
- 16. Singh A B, Chatarji. M, Singh B P and Gangal S V. (1990). Air borne viable fungi in Library before and after agitation of books, *Indian Journ. Aerobiol.* 3 (1&2), pp 32-38.
- 17. Tilak, S.T.(1989). Air borne pollen and fungal spores, Vaijanti Prakashan, Aurangabad
- Tilak S T, Pillai S G and Saibaba (1982). Component of aerospora inside the library and its relevance to book deterioration. Proc. I st Nat. Conf. Env. Biol. pp-173-177.
- 19. Tiwari K L, Jadhav S K and Saluja P .(2004). Studies on aeromycoflora of library, Botany Department and Garden, *Research Link*, Vol.14: III (4), Indore.
- Tripathi S N .(1987). Fungal aerospora inside the central library of Gorakhpur University, Proceedings, Water, air and soil pollution, 34, 125-134.
- 21. Upadhyay, H and A.K. Sahu. (2005). Fungal flora inside the library environment, Gwalior University, *Indian Journal of Aerobiology*. Vol 18, pp 6-11.
- 22. Verma K S and Khare K. (1987). Fungal spores which damage paper material in library, Atmospheric bio pollution. Ed. N Chandra pp193-197.
- 23. Vittal B P R and Glori A L .(1985). Air borne fungal spores of a library in India, Grana, 24: 129-