
Rointh Sniugia iu bias unimerizazu 1951

BIONANO FRONTIER

Vol.8 Issue - 1

January- June 2015



A Peer Reviewed / Referred Journal
of Science & Technology

Published From Wumbai www.bionononienag

SEASONAL FLUCTUATION OF AIR BORNE FUNGLIN THE COLLEGE

PATEL S.L.

Arts, Science and Commerce College, Ozar (MIG), Nashik, India 422 206, drsipatel a hotmail.com

Received on:10.10.2014 and Accepted Revised on:30.12.2014.

ABSTRACT: The present study was undertaken to identify and to find out the concentration of the various aero allergenic fungal spores in the indoor environment of the College library. The mycoflora of indoor and outdoor environment of library in Ozar (MIG) College was investigated for the period of one year from October 2013 to September 2014 by Humedia Petri plate air sampler No. LA-030 with Himedia readymade Rose Bengal Chloramphinicol Agar culture plates techniques. There were total 21 air borne fungal colonies were isolated and identified in the numbient air of college library. Aspergitus (19.82% and 28.2%), Cladosparium (8.99% and 7.20%) Penicillium 8.23% and 5.30%). Fusarium (5.34% and 4.50%) and Alternaria (6.10% and 9.50%) were found to be all time occurrence and shown predominance in the outside and inside the library environment respectively. Lignocellulosic degrading fungus like Chaetomium (5.18% and 4.19%) was also recorded in high humid and low temperature conditions, Alternaria, Fusarium Helminthosporium, Trichoderma, Rhizopus, Curvularia, Tetraplaa, Periconia, Stemphyllum and some unidentified colonies were recorded in high concentration during the investigation with little variation in their concentration. Repeated exposure to the library environment causes some health hazards like sneezing, watery eyes, coughing and uneasy feelings in some sensitive individuals which is in relation with the meteorological parameters and concentration of allergic spores present in the ambient air of the library.

Keywords: Library, Allergy, Airborne fungi, weather conditions

INTRODUCTION:

Today, more than 300 million of the world population is known to suffer from one or other allergic ailments affecting the socioeconomic quality of the life. The respiratory allergy is the most severe disorder including asthma, rhinitis, dermatitis, hay fever, skin rashes, sneezing, coughing, running nose, watering eyes etc. The major causative agents for the respiratory allergy are pollen grains, fungal spores, dust, mites, insect debris etc (Pande 2012).

Libraries are the main information and knowledge centres for the educational institutes. Due to suspended air borne particles in the library environment, both the readers and books get infected at more or less percentage. The enclosed environment of the library offer unique substrate such as binding glue, paper and other organic material which support the active fungal growth. Because of light weight and microscopic nature, the spores of fungi remain suspended in the air for long period. The concentration of these bioparticles shows great variation from time, season, altitude and weather conditions (Lacey 1991). Since the study of allergic aerospora of library environment is of great importance and continuous exposure to the library causes the respiratory disorders. A very meagre studies were reported on similar matter (Burge 1978, Vittal 1985, Tripathi 1987, Singh 1990, Agashe 1996, Ghani 1993, Verma 1996. Tilak 1997 Saoji 1997. Atluri 2002, Md. Shabbir 2003. Upadhyay 2005, Giri 2012,) hence the present work is undertaken to study the allergic air borne fungal spores in the ambient air of library.

MATERIALS AND METHODS:

The town Ozar (MIG), Maharashtra, is well known for its grape and onion production and similarly for oir craft production at Hindustan Aeronauties Limited. The arts,

Science and Commerce College, Ozar (MIG) was established along with central library in the year 1985. The total carpet area of the library is 1901 sq. fts consisting total 24,130 valuable books including text, reference books, periodicals and journals worth of Rs.27.38018/- up to year March 2014. The books were arranged subject wise and kept on open and closed steel racks keeping 3" distance between the racks.

The air sampling in the College library was done by using Himedia air sampler No. 030 with readymade Himedia Rose Bengal Chloramphinicol nutrient media plates. The air sampler was loaded with nutrient media plates and placed on the reading table for 10 minutes every forth night between 11.00 am to 11.30 am. i. e during peak hours of book transactions. The sampling of the air was done outside and inside the library environment from the months of October 2013 to March 2014 in the first season and April to September 2014 in the second season. After the exposure, the petriplates are brought to the botany laboratory for the incubation at the temperature 25-27 °C for 4-5 days. The Colony Forming Units (CFU) were counted and Total Fungal Counts (TFC) were expressed as Colony Forming Units per cubic meter of air (CFUs/M3)

The fungal colonies detected per unit volume of air is calculated as under

CFUs/M3= Number of Colonies on nutrient medium plates (NC)X 25

Sampling time in minus (T)

After the incubation period of 5-6 days the colonies were appeared on the nutrient medium petriplates. These colonies were counted and identified up to genus level with the help of available literature (Barnett 1971, Tilak 1985)

05

Print ISSN 0974-0678. Online: 2320-9593. www.bionanofrontier.org

BIONANO FRONTIER Vol. 8 (1) January-June 2015

RESULTS AND DISCUSSIONS

Studies reveals the total 21 types of fungal colonies recorded in the ambient air of the library. High concentrations of fungal colonies were recorded in the library environment (326) than outside the library. Highest concentration of colonies was recorded to be of Aspergillus (inside 28.57 % and outside 17.485%). Cladosporium (13.19% and 12.14 %). Penicillium (8.9% and 4.29%). Helminthosporium (6.13 % and 5.0%) and Alternaria (4.29% and 7.86%) to the total colonies recorded inside and the outside the library environment respectively (Table 01 and 02)

In addition to the cellulose, the paper also contains the lignin, hemicellulose, pectin, waxes, tannins and minerals useful for the fungal growth. Beside the impurities formed during the paper production are helpful for the nutrition of fungi (Tilak S T 1985). The cellulytic activity of some paper deteriorating fungi produces endogluconase enzymes which damages the paper quality of the library book (Verma K S 1987) indicate the role of fungi in book deterioration.

Present studies indicates the high concentration of the fungal colonies outside the library in the months of October to March 2013 while there was gradual increase the concentration inside the library from the Months of April to September 2014 might be due to high humid and low temperature conditions. (Fig.01 & 02). Some fungal types like *Drechslera, Epicoccum*, and *Geotrichum* were only reported in the rainy months of July to September 2014. High concentration of fungal colonies (713) inside the library environment indicates humid and damp environmental conditions which would be favourable for the growth of moulds on the books. Chauhan et al (2006) reported the similar observation by studying 7 different libraries at Agra. The adverse effects of fungal components not only deteriorate the paper material but also affect the library staff.


Agashe (1991) also reported 23 types of fungal colonies in the indoor environment of library at Bengaluru. Giri S K (2012)

reported 36 fungal spore types from the ambient air of library at Nagpur adding highest concentration of Aspergillus, Chadosporium, Alternaria and Rhizopus spores. Predominence of Aspergillus and Chadosporium in the present studies is in relation with the reports of Vittal (1985) and Singh (1990). The cellulose degrading fungi like Chaetomium was also recorded (5.48% and 4.19%) indicating the threat to the stored books in the library environment. Dhawan (2005) critically reviewed the role and aerobiodeteriogens on paper and museum material indicating high concentration of versatile lungi like Aspergillus, Peniculium, Cladosporium and Fusarium.

As indicated by Nandimuthan (1995) and Tripathi (1987) the present studies also reveals high concentration of fungal colonies during the high humid, low temperature conditions in the month of December 2013 and January 2014 than in high temperature and low humid conditions in the month of February and March 2014. Since the study of allergic airospora of library environment is of great importance and continuous exposure to the library causes the respiratory disorders. A very meagre studies were reported on similar matter (Tilak S T (1982), Verma K S (1987), Tripathi S N (1987), Nandimuthan (1995), Shaney (2001), Sharma (2004), Tiwari K L (2004). And Mujumdar (2005). The studies inside and outside the library environment is of great importance since the library environment is loaded with the biopollutants that can be harmful to the health of the library users, beside reducing the life of books (Agashe S N 1996)

ACKNOWLEDGEMENTS

The author is extremely thankful to The Director, BCUD, Savitribai Phule Pune University, Pune for the financial assistance and The Principal Dr. A.P.Patil, MVP Samaj's Arts, Science and Commerce College, Ozar (MIG) for permitting air sampling in the College library and providing laboratory facilities.



Ing. 01 Sumber of Copy (all or consider yet the 2-septime), pulsed and mode in a college from a time of consent of the consent

Table: 01

Sr No	Fungal Types	Number of colonies recorded on the petriplates (Outside the library)														
		Oct		Dec		1	1		May		Jul	Ang	Sep	Total	-	CFU/M
1	Alternaria	3	4	1	2	1	3	-	2	4	7	8	5	40	6.10	100
2	Aspergillus	12	7	11	12	7	8	6	8	5	15	22	17	130	19.82	325
3	Cercospora	3	2	2	-		2			3	-	5	-	17	2.59	42.5
4	Chaetomium	3	5	4	6	3	2	3		3	-	2	3	34	5.18	85
5	Cladosporium	12	3	10	11	4	3	2	4	5	-	3	2	59	8,99	147.5
6	Corynospora	-	-	-	-		-	2	7		5	2	1	17	2.59	42.5
7	Curvularia	-	-	3	4	1		-	2	5		5		20	3.05	50
8	Drechslera	-	-	-	-		-	-	-	-		-		-	-	-
9	Ерісоссит	2	1	-	2	I	- 1	-	-	-	3	7	-	17	2.59	42.5
10	Fusarium	3	2	4	.3	5	4	2	1	2	-1	3	2	35	5.34	87.5
11	Geotrichum	2	-	3	-	4	6	-	- 1		-		-	15	2.29	37.5
12	Helminthosporium	2	4	6	4	3	1	-	3	1		3	3	3()	4.57	75
13	Nigrospora	-	-	1	1	2	+	-		3		2	2	11	1.68	27.5
14	Penicillium	3	5	7	8	4	2	2	3	.3	7	7	3	54	8.23	135
15	Phoma	-	4	-	-	-	-	-	-			-	-	1	0.15	2.5
16	Periconia	-	1	2	1	1	-	-		6	3	2	5	21	3.20	52.5
17	Pringsheimia	-	-	-	1	1	-	-			- 1	-	2	5	0.76	12.5
18	Rhizopus	3	5	2	3	1	-	-	-	2	3	7	3	29	4.42	72.5
19	Stemphyllum	2	5	3	7	2	1		1	3	6	-	4	34	5.18	85
20	Trichoderma	3	4	2	2	4	5	2		2	2	1	2	29	4.42	72.5
21	Unclassified	4	5	7	4	2	7	4	3	3	7	3	ŋ	58	8.84	145
	Total	57	54	68	71	46	45	23	34	50	63	82	63	656	100	1640

Table:02

Sr No	Fungal Types	Number of colonies recorded on the petriplates (Inside the library)														
		Oct	Nov	Dec	Y		1		May		Jul	Aug	Sep	Total		
1	Alternaria	2	.3	2	2	-	2	3	-	23	12	9	10	68	9.54	170
2	Aspergillus	3	3	7	10	12	5	2	17	33	42	23	44	201	28.19	502.5
3	Cercospora	-	-	2	2	1	1	3	-	.5	4	2	7	27	3.79	67.5
4	Chaetomium	-	-	-	2	1	1	4	-	7	5	7	3	30	4.21	75
5	Cladosporium	2	4	5	3	2	1	9	7	2	5	7	4	51	7.15	127.5
6	Corynospora	-	-	*	-		-	3	2	4	-	7	1	16	2.24	40
7	Curvularia	-	-	-	-	2	1	3	- 1	4	-	7	7	24	3.37	60
8	Drechslera	- 1	1		-	-	-	-	-	3	5	2	2	12	1.68	30
9	Ерісоссит	-	-	-	2	-	-	2		1	4	-		. ()	1.26	22.5
10	Fusarium -	2	1	-	-		-	4	2	4	8	7	4	32	4.49	80
11	Geotrichum	-		-	-	-	-	2	-	3		4	6	15	2.10	37.5
12	Helminthosporium	-	1	2	1	2	1	2		3	5	3	8	28	3.93	70
13	Nigrospora			2	1	1	-	-	3			2		0	1.26	22.5
14	Penicillium	3	2	1	- 1	-	-	2	7	3	7	7	6	38	5.33	95
15	Phoma	-		-	-	- 1	- 1		1	-	7	3	-	11	1.54	27.5
16	Periconia	-	-	-	1	1	-	1	2	1	4	3	2	15	2.10	37.5
17	Pringsheimia	-	1	-	-	-	- 1	2	- 1		1	-	-	4	0.56	10
18	Rhizopus	2	1	1	-	2	2	1	2	5	3	7	7	3.3	4.63	82.5
19	Stemphyllum	2	1	3	1				-	3	4	1	2	17	2.38	42.5
20	Trichoderma	2	1	1	-				2		7	4	3	20	2.81	50
21	Unclassified	3	4	2	1		4	3	3	4	10	12	6	53	7.43	132.5
	Total	21	22	28	26	25	18	46	48	108	133	117	121	713	100	1782.5

REFERENCES

- Agashe, S.N. and Anuradha. (1996). Aeromycological studies of a library in Bangalore city. Ind. Journ. Aerobiol. Vol. 11: 24-28.
- Atluri, J.B and Padmini. (2002). Aeromycoflora of Andra university main library. *Indian Journal of Aerohology*. Vol 18 (II), pp.47-58.
- Barnett, H.L. and Hunter. B.B. (1972). Illustrated genera of imperfect fungi. Burgess. Co. Minneapolis, Minnesota, U.S.A.
- Burge H A. (1991). Bioaerosols: prevention and health effects in the indoor environment. *Journ. Allergy clinical* immunology, vol (26) 05, 687-701
- Chauhan, S.V.S. A Kulkarni and Goyal (2006). Air borne fungal spores of Agra city with special reference to allergic significance, *Indian Journal of Terobiology*. Vol 17, pp 17-22.
- 6) Dhavan Sashi, and Neet Nigam. (2005). Fungal diversity of paper material and its role in bio deterioration in fungal diversity and biotechnology. ALK Ikai and S.K. Deshmukh Scientific publisher Dodhpur. India PP 93-113.
- Ghani, T and Shanta Kale. (1993). Aeromycoflora inside and outside the College library. *Indian Journal of Environ Health*. Vol 3 (II), pp 1-5.
- Lacey, J. (1991). Aerobiology and health. Role of aerobiology Fungal spores in respiratory diseases in. Oxford and IPH publishing Co. Ltd, New Delhi
- Mohammad Shabbir, S. Nicky, Johnson and Rajasab. (2003). Acromycoflora of Gujarat University library. Indian Journal of Aerobiology. Vol 16, pp 28-32.
- Mujumdar M R and Surendra Hazare. (2005). Assessment of fungal contaminents in library of Kolkata, Ind. Journ. Aerobiol. 13: 398-403.
- Nandimuthan and Vittal B P R. (1995). A volumetric survey of culturable moulds in Library. Ind. Journ.

- derubial & 12-18
- Pande B.N. (2012). Bioacrosofs as affergens. Bionano Frontier, Vol. 5 (2), 164-160.
- Saoji A A and Gift S K. (1997). Concentration of aero allergenic fungal spaces in intramural environment of Nagpur City-Hospital ward and library. 5° International conference on Actiobiology pp-211-218.
- 14) Shaney, Manju and Purwar (2001). Aeromycoflora inside a library of Affahabad University, Ind. Journ. Aerobiol, 14, 20-23.
- Sharma K B and Basumaitrai (2004). Air borne fungal speces in library and their association with paper and leather material, Incl. Journ. Aerobiol., 14: 35-43.
- 16) Singh A B. Chatarji, M, Singh B P and Gangal S V. (1990). Air borne viable fingi in Library before and after agitation of books. *Inclian Journ. Aerobiol.* 3 (1&2), pp. 32-38.
- Tilak, S.T.(1989). Air borne pollen and fungal spores, Varjanti Prakashan, Aurangahad
- 18) Tilak S T, Pilfai S G and Saibaba (1982). Component of aerospora inside the library and its relevance to book deterioration. Proc. I" Nat. Conf. Em. Biol. pp-173-177.
- Tiwari K L, Jadhav S K and Saluja P. (2004). Studies on aeromycoflora of library. Botany Department and Garden, Research Link, Vol. 14: III (4). Indore.
- Iripathi S N. (1987). Fungal aerospora inside the central library of Gorakhpur University. Proceedings, Water, air and soil pollution, 34, 125-134.
- Upadhyay, H and A.K. Sahu. (2005). Fungal flora inside the library environment, Gwalior University. *Indian Journal of Aerobiology*. Vol 18, pp 6-11.
- Verma K S and Khare K. (1987). Fungal spores which damage paper material in library. Atmospheric bio pollution. Ed. N Chandra pp 193-197.
- Vittal B P R and Glori A L .(1985). Air borne fungal spores of a library in India. Grana. 24: 129-132.